

Observing and modelling Earth's albedo and its interhemispheric symmetry

Frida Bender, Aiden Jönsson

Earth Radiation Budget Worskhop, Oct 13 2022



SMHI

Bolin Centre for Climate Research



Why albedo?



visibleearth.nasa.gov

Bolin Centre for Climate Research

A collaboration between Stockholm University, KTH and the Swedish Meteorological and Hydrological Institute

Why albedo?

- $S_0/4(1 - \alpha) = \sigma T^4$



visibleearth.nasa.gov

Bolin Centre for Climate Research

A collaboration between Stockholm University, KTH and the Swedish Meteorological and Hydrological Institute

Why albedo?

- $S_0/4(1 - \alpha) = \sigma T^4$
- $\alpha = f(\text{cloud})$



visibleearth.nasa.gov

Bolin Centre for Climate Research

A collaboration between Stockholm University, KTH and the Swedish Meteorological and Hydrological Institute

Why albedo?

- $S_0/4(1 - \alpha) = \sigma T^4$
- $\alpha = f(\text{cloud})$
- $\alpha(\text{NH}) \approx \alpha(\text{SH})$




visibleearth.nasa.gov

Bolin Centre for Climate Research

A collaboration between Stockholm University, KTH and the Swedish Meteorological and Hydrological Institute

Why care about cloud representation in global models?




-Multiple lines of evidence narrows ECS uncertainty range in AR6

-ECS best estimate 3C and *very likely* range 2C to 5C

-Cloud feedback dominant source of uncertainty in future projections


IPCC AR6 Fig 7.18 (cropped)

Cloud feedback in CMIP5 and CMIP6



- More positive cloud feedback in CMIP6 than in CMIP5 explains higher sensitivity

Global albedo – CMIP3




Bolin Centre for Climate Research


A collaboration between Stockholm University, KTH and the Swedish Meteorological and Hydrological Institute

Bender et al (2006), Tellus


Model clouds overall "Too few, too bright"



"Too few, too bright" remains in CMIP6



Interhemispheric albedo symmetry




Bolin Centre for Climate Research


A collaboration between Stockholm University, KTH and the Swedish Meteorological and Hydrological Institute

E.g. Voigt et al. (2013, 2014), Stephens et al. (2015), Bender et al. (2017),
Datseris & Stevens (2021), Jönsson & Bender (2022), Diamond et al. (2022)


Regional asymmetry => hemispheric symmetry (CMIP5)




Regional asymmetry => hemispheric symmetry (CMIP5)




Regional asymmetry => hemispheric symmetry (CMIP5)




Regional asymmetry => hemispheric symmetry (CMIP5)



Regional asymmetry => hemispheric symmetry (CMIP5)



Regional asymmetry => hemispheric symmetry (CMIP6)




Bolin Centre for Climate Research

A collaboration between Stockholm University, KTH and the Swedish Meteorological and Hydrological Institute


Courtesy Marcus Karlsson

Albedo symmetry in CERES EBAF

- Inter-hemispheric differences in reflected SW radiation within 0.1 Wm^{-2}
- Greater reflection from clouds in SH balances greater surface albedo and clear-sky albedo in NH




Patterns of anomalous asymmetry composites



- The variability in asymmetry is strongly determined by tropical and subtropical clouds, particularly those related to non-neutral phases of ENSO.


Albedo symmetry in CMIP6 models



- CMIP6 models underestimate symmetry, and overestimate variability
- Fixed SST brings models closer to symmetry
- Model bias in symmetry driven by midlatitude clouds

Asymmetry response to strong forcing

- Models lose NH sea ice in response to strong forcing
- Some models regain symmetry
- Restoration can be related to SH cloud loss, or to NH cloud gain - with different implications for cloud feedback







Asymmetry response to strong forcing

Models that restore more...

...lose more clouds in the SH mid-latitudes...

...and are typically at the high-ECS end




Bolin Centre for Climate Research

A collaboration between Stockholm University, KTH and the Swedish Meteorological and Hydrological Institute


Courtesy Aiden Jönsson

Factors controlling SH midlatitude cloud



- Models with greater drop in surface wind speed, more SST increase, more decrease in equator-to-pole gradient lose more cloud
- Similar cloud controlling factors determine present-day variability


Albedo symmetry in storm resolving models



	NH-SH mean
ICON ngc2009 (5 km)	-0.89%
ICON ngc2012 (10 km)	-1.4%
CERES EBAF (clim)	-0.03%

- Storm-resolving (5km, 10km) ICON reproduces interhemispheric albedo symmetry well
- ICON (5km) represents absolute albedo well

Albedo symmetry in storm resolving models



- Increasing resolution from 10 km to 5km reduces (the positive) CRE bias in SH midlatitude storm tracks
- Storm track position moves poleward with increasing resolution, in better agreement with CERES

Summary

- Cloud-compensation of clear-sky asymmetry stems from mid-latitude cloud amount and cloud albedo, and subtropical cloud amount
- Midlatitude cloud properties are determining of level of symmetry, and model bias, while tropical cloud distribution determines its variability
- Restoration from CO₂-induced imbalance is driven by SH mid-latitude clouds. Most restoration seen in most sensitive models.
- Storm-resolving models show promise of improved representation of albedo symmetry

Bolin Centre for Climate Research

A collaboration between Stockholm University, KTH and the Swedish Meteorological and Hydrological Institute

